

“ $\Omega > 1$ ”

“Sky-Notes” of the Open University Astronomy Club.

January 2022.

A Very Happy New Year and Best Wishes for 2022 to all readers!

Recent Events.

If you have any images and/or reports of recent events please contact Sheridan so that he can put them on the Club website.

If you wish to present them at a “Clubnight” meeting please contact Adrian or myself before the meeting starts.

Highlights of the Month.

4th **Earth at Perihelion.**
3rd Peak of **Quadrantids** meteor shower.
7th **Mercury at Greatest Elongation E.**
8th **Venus at Inferior Conjunction.**
16th **Pluto in Conjunction with the Sun.**
23rd **Mercury at Inferior Conjunction.**
Mercury. Low in the SW evening twilight.
Venus. Watch for its reappearance mid-month low in SE dawn twilight.
Mars. Low in SE predawn sky.
Jupiter. Low in the WSW early evening sky.
Saturn. Becoming lost low in the SW evening twilight by mid-month.
Uranus. Well placed in Aries for evening observation.
Neptune. An early evening object low in SW sky.
Asteroids at Opposition – see notes below.
Comets and Meteor Showers – see notes below.

Forthcoming Meetings.

OUAC.

The next **Clubnight** will be held on **Tuesday 11th January at 19:30** in the Activity Room of the Sports Pavilion. The programme will include short presentations by members, “Sky-Notes” and **most importantly** to discuss and formulate the programme for the remaining session in still difficult circumstances with respect to Covid.

Software.

A very useful item of Planetarium software is “**Stellarium**” and it’s **FREE!**
Go to the website and download it and the associated user manual.

1. The Solar system.

Note all times shown are UT.

Earth.

Perihelion 04^d 06^h 55^m 147,105,053 km.

Aurora.

Long hours of darkness improve the opportunity for observing potential aurora.

Keep tuned to the www.spaceweather.com site for updates.

Subscribe (free) to the UK AuroraWatch website to receive alerts.

ISS.

Completes a series of morning passes during the first week of the month. Commences a series of evening passes at the end of the third week of the month.

For this and other “bright” satellites go to the “Heavens Above” website and set-up for your location.

Alternatively go to “spaceweather” website and click the “Flybys” button and follow the instructions to set-up forecasts for your location.

Add to your “favourites”.

Sunrise and Sunset.

Bedford.

Latitude 52° 6.9’N Longitude 0° 28.1’W

Date.	Rise.	Transit.	Set.
01	08 ^h 11 ^m	12 ^h 06 ^m	16 ^h 00 ^m
08	08 ^h 08 ^m	12 ^h 09 ^m	16 ^h 09 ^m
15	08 ^h 04 ^m	12 ^h 11 ^m	16 ^h 19 ^m
22	07 ^h 56 ^m	12 ^h 14 ^m	16 ^h 31 ^m
29	07 ^h 47 ^m	12 ^h 15 ^m	16 ^h 44 ^m

Produced using Starry Night Pro v6.

The Sun.

To prevent permanent damage to your eyes avoid looking at the Sun directly and never with binoculars or a telescope unless special (expensive!) filters are used.

The safest way is the simplest – project the image of the Sun onto grey or white card.

Take additional care if your telescope has plastic components. Plastic melts!

If you have or have access to observe in h-alpha the rewards are much greater.

Solar Cycle 25 is well underway.

Keep in touch with the Solar Dynamics Observatory satellite at <http://sdo.gsfc.nasa.gov/>

Add the “spaceweather” and “Soho” websites to your “favourite” websites.

The Moon.

Phases:



Produced using Lunar Phase Pro.

New	02 ^d 18 ^h 33 ^m
First Quarter	09 ^d 18 ^h 11 ^m
Full	17 ^d 23 ^h 48 ^m
Last Quarter	25 ^d 13 ^h 41 ^m

Apsides:

Perigee	01 ^d 23 ^h	Diameter. 33' 52"	Distance. 358,032km.
Apogee	14 ^d 09 ^h	Diameter. 29' 57"	Distance. 405,804km.
Perigee	30 ^d 07 ^h	Diameter. 33' 29"	Distance. 362,253km.

Observing, Imaging Opportunities and Challenges.

For northern observers:

The waxing crescent Moon is becoming better placed.

The waxing gibbous Moon is very well placed.

The Full Moon is very well placed.

The waning gibbous Moon is well placed.

The waning crescent Moon is less well placed.

The Moon cont.

Observe the regions along the terminator (sunrise and sunset on the Moon) where the low angle of the Sun highlights lunar topography. A basic lunar map is all you need to get started. *Sky & Telescopes* "Lunar 100 Card" is another good starting point. If you are starting out on photography and/or imaging the Moon provides an excellent target.

On 1st (difficult!), 30th and 31st (difficult!) try locating the very thin crescent Moon very low in the ESE dawn skies **before sunrise**.

On 3rd (difficult!) and 4th try locating the very thin crescent Moon in SW evening twilight **after sunset**.

If you can take images of the above so much the better.

Lunar Occultations.

Unlike the gradual disappearance of a planet (small disc) a star vanishes instantly demonstrating that it is a point source of light as viewed from the earth. For all occultation events start observing 10 to 15 minutes before the predicted time to identify the required star and to allow for slightly different time if you are not at Greenwich. Use an accurate watch to record the time that *you* observe the occultation remembering that times are UT not BST. Enter details in your observing log.

Further details of occultations can be found in current *BAA Handbook* and monthly periodicals such as *Astronomy Now* and *Sky at Night*.

On the morning of 26th the double star **alpha (α) Libra = Zubenelgenubi (+2.8)** is occulted by the Moon.

Disappearance behind the bright limb (DB) at 05^h 37.2^m.

Reappearance from behind the dark limb (RB) at 06^h 48.1^m.

Times shown are for Greenwich so start your observations earlier, say 15 to 20 mins.

Useful article in January issue of *Astronomy Now*.

The Planets.

Mercury.

Low in the SW evening twilight in the first half of the month.

Try spotting it on the 4th when it forms a pleasant triangle with Saturn and a thin crescent Moon. Higher and to the east is Jupiter.

Greatest Elongation E (19°) on 7th.

Inferior Conjunction on 23rd.

Moon close on 4th and 30th.

Date.	Mag.	Dia.	Phase.	Rise.	Transit.	Set.
01	-0.7	5.9''	0.77	09 ^h 24 ^m	13 ^h 24 ^m	17 ^h 23 ^m
07	-0.5	6.8''	0.59	09 ^h 14 ^m	13 ^h 30 ^m	17 ^h 46 ^m
31	+1.4	9.5''	0.18	06 ^h 40 ^m	11 ^h 06 ^m	15 ^h 31 ^m

Venus.

May just be spotted very low in the SW evening twilight in the first two days of the month. **Don't sweep with binoculars until the Sun has completely set!**

Inferior Conjunction on 8th.

Watch for its reappearance low in the SE morning twilight mid-month.

Moon close on 3rd and 30th.

Date.	Mag.	Dia.	Phase.	Rise.	Transit.	Set.
01	-4.3	60''	0.02	08 ^h 33 ^m	12 ^h 55 ^m	17 ^h 16 ^m
15	-4.2	60''	0.02	06 ^h 51 ^m	11 ^h 25 ^m	15 ^h 57 ^m
31	-4.6	50''	0.15	05 ^h 30 ^m	10 ^h 06 ^m	14 ^h 41 ^m

Mars.

Low in ESE predawn sky.

Moon close on 29th.

Date.	Mag.	Dia.	Phase.	Rise.	Transit.	Set.
01	+1.5	4.0''	0.98	06 ^h 10 ^m	10 ^h 06 ^m	14 ^h 01 ^m
31	+1.4	4.3''	0.96	05 ^h 55 ^m	09 ^h 41 ^m	13 ^h 28 ^m

Jupiter.

Low in the WSW evening sky.

Moon close on 5th and 6th.

See BAA *Handbook* and/or monthly periodicals for satellite phenomena.

Date.	Mag.	Dia.	Rise.	Transit.	Set.
01	-2.1	35''	10 ^h 30 ^m	15 ^h 29 ^m	20 ^h 28 ^m
31	-2.0	34''	08 ^h 44 ^m	13 ^h 55 ^m	19 ^h 08 ^m

Saturn.

Becoming lost low in the SW evening twilight by mid-month.

Try spotting it on the 4th when it forms a pleasant triangle with Mercury and a thin crescent Moon. Higher and to the east is Jupiter.

Conjunction with the Sun on February 4th.

Moon close on 4th.

Date.	Mag.	Dia.	Rise.	Transit.	Set.
01	+0.7	15''	09 ^h 51 ^m	14 ^h 16 ^m	18 ^h 41 ^m
15	+0.7	15''	08 ^h 59 ^m	13 ^h 27 ^m	17 ^h 55 ^m

Uranus.

Located in Aries and well placed for evening observation.

Moon close 11th.

Date.	Mag.	Dia.	Rise.	Transit.	Set.
01	+5.7	3.7''	12 ^h 29 ^m	19 ^h 51 ^m	03 ^h 12 ^m
31	+5.8	3.6''	10 ^h 31 ^m	17 ^h 53 ^m	01 ^h 18 ^m

Neptune.

Sinking into the SW early evening skies.

Moon close 7th.

Date.	Mag.	Dia.	Rise.	Transit.	Set.
01	+7.9	2.3''	11 ^h 06 ^m	16 ^h 44 ^m	22 ^h 23 ^m
31	+7.9	2.2''	09 ^h 09 ^m	14 ^h 49 ^m	20 ^h 29 ^m

Dwarf Planets.

Ceres.

A 8th mag object in Taurus. Moon close on 12th.

Eris.

A CCD target object in Cetus.

Haumea.

A CCD target located in Bootes.

MakeMake.

A CCD target in Coma Berenices.

Pluto.

Not observable. **Conjunction** with the Sun on 16th.

Asteroids. (Approx mag +10.5 or brighter).

Vesta (4).

An 8th mag object located in Sagittarius emerging into the predawn SE sky at the end of the month..

Iris (7).

Located in Gemini. Mag +7.7 at **Opposition** on 13th.

Charts and details of asteroids one month either side of opposition are available at:

http://britastro.org/computing/charts_asteroid.html

See also the "Heavens above" website, the *BAA Handbook* and/or monthly periodicals.

Comets.

A good month for moderately bright comets!

C/2019 L3 (ATLAS). A 9th mag object tracking through Gemini.

19P/Borrelly. A 8th/9th mag object tracking through Cetus into Pisces at the end of the month..

67P/Churyumov-Gerasimenko. A 9th mag object tracking through NvCancer.

Charts and details of selected comets are available at:

http://britastro.org/computing/charts_comet.html

See also the “Heavens above” website, the *BAA Handbook* and/or monthly periodicals.

Meteor Showers.

The **Quadrantids** are active from 1st – 6th January with peak activity on 3^d 17^h, ZHR = 80+. Rather narrow peak but **very favourable as Moonlight does not interfere!**

There are always **Sporadic** events and the chance of a brilliant fireball. The latter should be recorded and reported.

Near Earth Objects.

Please refer to www.spaceweather.com for updates.

Eclipses.

No Lunar or Solar Eclipses this month.

2. The Deep Sky.

Abbreviations used.

M = Messier object (Shown in **bold**).

NGC = New General Catalogue. IC = Index Catalogue (Extension of the NGC).

ds = double star. ts = triple star. ms = multiple star. vs = variable star.

gc = globular cluster. oc = open cluster. pn = planetary nebula.

en = emission nebula. rn = reflection nebula. sg = spiral galaxy.

eg = elliptical galaxy. lg = lenticular galaxy. ir = irregular galaxy.

pg = peculiar galaxy. snr = super nova remnant. ly = light year.

The magnitude of an object, excluding double, triple, multiple and variable stars, is shown in brackets e.g. (6.5).

All magnitudes are + unless otherwise shown.

2.1 Variable Stars of the month.

Beta (β) Persei, Algol.

Range +2.2 to +3.4, period 2.7 days. Becoming well placed by mid evening.

Favourable minima at “social hours” occur on 2^d 23.5^h, 5^d 20.3^h, 25^d 22.0^h and 28^d 18.8^h.

Delta (δ) Cephei.

Range 3.5 to 4.4, period 5.37 days. The prototype for the Cepheid class of variable stars.

Their period-luminosity relationship has led them to being used as “standard candles” in measuring distances to nearby galaxies.

Mu (μ) Cephei.

Range 3.7 to 5.0, approximate period 755 days. A semi-regular variable star famous for its striking red colour being fittingly called “Herschel’s Garnet Star”. It is the reddest naked eye star visible from the northern hemisphere. Its colour may show signs of variability.

U Ori.

Well placed for nightlong observation. Brightening from minimum (+12) in December 2021 towards maximum (+6.3) in May.

2.2 Double Stars of the month.

Beta Cam. See notes below.

Alpha Gem (Castor). See notes below.

Delta Gem. See notes below.

Gamma Lep. See notes below.

Beta Mon. See notes below.

Beta Ori (Rigel). See notes below.

Sigma Ori. See notes below.

Theta-1 Ori (The Trapezeium). See notes below.

Struve 742 Tau. See notes below.

2.3 This Month's Constellations - Double Stars/Star Clusters/Nebulae/Galaxies.

Auriga (Aur).

NGC1960 (M36) (6.0) oc. Large bright grouping. In same low power field as M38.

NGC2099 (M37) (5.6) oc. Richest and brightest of the three Messier star clusters in Auriga with 150 stars brighter than 12th magnitude.

NGC1912 (M38) (6.4) oc. Larger than M36. Many bright stars arranged in pairs.

The above are excellent objects for photography. Guided exposures of a few minutes will be necessary. CCD images require much shorter exposures.

NGC1664 (7.6) oc. Fine cluster on the border of Auriga and Perseus.

NGC1778 (7.7) oc. A 6" telescope will show about 20 stars. Larger apertures will reveal more.

NGC1857 (7.0) oc. Hazy patch surrounding an orange 7th magnitude star which interferes with viewing the fainter stars.

NGC1893 (7.5) oc. Fine, though rather sparse cluster. 8"+ telescopes under dark skies may begin to reveal the pale light of the brightest part of the emission nebula IC410 which pervades the star cluster.

NGC1907 (8.2) oc. This small cluster lies just west of M38 appearing as a small smudge of light.

NGC1931 (11.3) en. An 8" telescopes from dark skies should reveal this small pea-nut shaped emission nebula.

NGC 2192 (10.9) oc. Not an easy object probably requiring a 6" telescope to locate and 10"+ to resolve.

NGC2281 (5.4) oc. Handful of stars often overlooked.

IC405 en (6v) The "Flaming Star Nebula". Illuminated by the star AE Aur which is a "runaway star" whose path can be traced back to Orion. At present the star is passing by/through the normally dark dust and gas cloud of IC405 and thus illuminating it. In the future as AE moves away the nebula will again become dark.

Camelopardalis (Cam).

Beta (β) ds. (4.0/8.6, sep. 80.8"). Wide yellow and blue pair (AB). From dark sites the system can be seen to be embedded in a streak of dark nebulosity. A third magnitude 11.8", sep (BC) 14.8", component is visible in moderate apertures.

1 ds. (5.7/6.8, sep. 10.3"). White and pale blue pair.

11 & 12 ds. (5.4/6.5, sep, 108.5") White pair. Easy object for binoculars and small telescopes.

29 ds. 6.5/9.5, sep. 25.1". Yellow primary with fainter pale blue secondary.

Struve (Σ)390 ds. 5.1/9.5, sep. 14.8". White primary with fainter purple secondary.

Struve (Σ)1122 ds. (7.8/7.8, sep. 15.4"). Fine pair of equally bright white stars.

Struve (Σ)1625 ds. 7.3/7.8, sep. 14.4". Almost equal pair of yellowish stars.

Struve (Σ)1694 ds. 5.3/5.8, sep.21.6". Easy pair of whitish stars.

NGC1501 (11.5) pn. Bright, large with a blue tint located about 2 degrees south of the open star cluster NGC1502. Resembles the "Eskimo Nebula" in Gemini. NGC1502 (5.7) oc. A fine open cluster. Extending NW of this cluster is a line of 9th and 10th magnitude stars that form "Kemble's Cascade". A beautiful sight in binoculars or a low power wide field eyepiece on small telescopes.

NGC2403 (8.4) sg. Visible in large binoculars. It lies at a distance of 8m light years and is possibly a member of the M81/M82 group.

IC342 (12.0) sg. Once considered to be a member of the Local Group of galaxies. Its low surface brightness makes it a challenge for moderate apertures. This face on spiral galaxy would be an impressive object if it were further away from the plane of the Milky Way.

Gemini (Gem).

Alpha (α) **Castor** ms. 1.9/2.9 sep 4.0". Close visual pair. However each of these is a spectroscopic binary. A more distant ninth magnitude star (red) forms part of an eclipsing binary system. A fascinating family!

Delta (δ) ds. 3.5/8.2 sep 5.8". Yellow primary with bluish secondary.

Kappa (κ) ds. 3.6/8.1 sep 7.1". Orange-yellow primary with bluish companion.

Lambda (λ) ds. 3.6/10.7 sep 9.6". Blue-white primary with bluish companion.

Σ 1108 (Struve) ds. 6.6/8.3 sep 11..5". Yellow primary with bluish companion.

M35 (5.1) oc. Just visible to the naked eye from dark sites. It is a superb object in telescopes. On its western edge lies the more distant open star cluster IC2158.

NGC2129 (10.2) oc. Located about a degree SW of IC2158.

NGC2266 (9.5) oc. Located about two degrees north of ϵ Gem.

NGC2392 (10.5) pn. The "Eskimo nebula" is a fine planetary nebula located about two degrees SE of δ . The nickname is derived from the appearance of a face surrounded by the hood of a parka.

NGC2420 (8.3) oc. Located about two degrees east of the "Eskimo".

Complete this deep-sky tour of Gemini by locating the open star clusters NGC2355 (9.7) and NGC2395 (7.1).

Lepus (Lep).

Lying beneath Orion Lepus is easily recognized by a quadrilateral of four third magnitude stars and contains a variety of deep-sky objects including one Messier object.

Alpha (α) ds; (2.6,11.1; sep. 35.8").

Beta (β) ds; (3.0/7.5; sep. 2.3").

Gamma (γ) ds. (3.7/6.3, sep. 96.3"). Fine yellow and pale orange pair.

Kappa (κ) ds; (4.5/7.4; sep. 2.6"). White and blue companions.

Iota (ι) ds; (4.5/10.8; sep. 12.7").

NGC1974 sg (11.8). Seen almost edge on.

NGC1904 (**M79**) gc (8.0). A fine globular cluster visible as a fuzzy spot in binoculars. Outer edges begin to resolve in 12" (30cm) telescopes.

IC418 pn (10.7). Very small but bright. Central 10.7 mag star surrounded by pale ring. Use a UHC or OIII filter for best results.

Monoceros (Mon).

This faint and rather indistinct constellation is located between Orion and Canis Minor.

Beta (β) ts. 4.7/5.2/6.1 Sep. AB = 7.3', sep BC = 2.8". Striking triple of bluish white stars.

Epsilon (ϵ) ts. 4.5/6.5 sep. 13.4". Close pair of pale yellow stars. The third mag 12.7 bluish white member is visible in 12"+ apertures.

NGC2244 oc (4.8). Fine open star cluster surrounded by NGC2237-9 "The Rosette Nebula" which is best seen using a UHC filter. Shows well in photographs.

NGC2261 en (10v). "Hubble's Variable Nebula". Located about 2° southwest of NGC2264 this a fascinating object and well worth monitoring for changes in shape and brightness due to the enveloped variable star R Monocerotis. The triangular wedge appears is almost comet like. Detailed star chart available for telescope owners.

NGC2264 oc + en (4.0) The "Christmas Tree Cluster". A fine open cluster with associated nebula that includes the "Cone Nebula".

NGC2323 (**M50**) oc (5.9). Superb open cluster.

There are many other open clusters in this area of the Milky Way - NGC's 2215(8.4), 2286(7.5), 2301(6.0), 2335(7.2), 2343(6.7), 2353(7.1) and 2506(7.6).

Orion (Ori).

This constellation dominates the winter skies and because it is so easily recognized forms one of the "key constellations" for finding other winter groupings.

Orion's two brightest stars provide a marked contrast. Betelgeuse is distinctly orange in colour. It is a red giant star entering old age. Rigel is a brilliant blue/white star indicating the exuberance of youth. Betelgeuse is slightly variable in brightness, range 0.1 - 0.9 and bears the designation α (alpha) indicating that it was brighter than Rigel, β (beta) (0.1), when stars were given these designations. Rigel is now the brighter of the two so either early magnitude estimates were wrong or Betelgeuse has dimmed slightly.

Beta (β) Rigel ds. 0.1/6.8 sep. 9.5". Brilliant bluish white primary with much fainter bluish secondary.

Eta (η) ds. 3.6/5.0 sep. 1.5". Close pair of white stars.

Delta (δ) ds. 2.0/6.9 sep. 52.6". Blue white primary with pale blue secondary.

Lambda (λ) ds. 3.5/5.6 sep. 4.4". White stars. Part of a quadruple system.

Theta-1 (θ) ms. "The Trapezium". AB: 6.7/7.9 sep. 8.8"; CD: 5.1/6.7 sep. 13.4". Superb object!

Iota (ι) ts. 2.8/7.3 sep. 11.3". White primary with pale blue secondary. The third reddish 11th magnitude component is located 50" away.

Sigma (σ) ms. 4.0/10.3 sep. 11.4. Colourful multiple star. Fainter triple star Σ 761 in same field.

Zeta (ζ) Alnitak ds. 1.9/4.0 sep. 2.3". Bluish white stars. Part of a triple system with the Flame Nebula (NGC2024) in the field of view.

NGC1976 (**M42**) en. One of the most famous objects in the sky. Marking Orion's sword the "Great Orion Nebula" is visible to the naked eye as a faint misty patch. A pair of binoculars or small telescope will begin to reveal detail. Increasing aperture and low power bring increasing rewards for the visual observer. Embedded in the nebula is **Theta (θ^1) Ori**. A group of four young stars, mags 5.4, 6.3, 6.8 and 7.0, aptly called "The Trapezium". The whole nebula is a stellar nursery with spectacular images being obtained from large Earth based telescopes and the Hubble Space Telescope. M42 is an ideal target for photography.

NGC1982 (**M43**) en. A small patch of nebulosity on the northern edge of M42.

NGC2024 (en), nicknamed "the Flame Nebula", surrounds ζ Ori.

IC434 en is a strip of nebulosity just south of ζ . The famous "Horse's Head Nebula" (Barnard 33) is a small dark intrusion seen dramatically in photographs. It provides one of the biggest challenges to visual observers requiring very dark transparent skies. Responds well to a H-beta nebula filter.

NGC2068 (**M78**) (8.0) rn is a small patch of nebulosity about two degrees NNE of ζ .

NGC2112 (9.1) oc is an open star cluster about two degrees east of M78.

Other open clusters worth locating are NGC2186 (8.7), NGC2169 (5.9) and NGC2175 (6.8) which superimposes a small patch of nebulosity NGC2174.

Long exposure photographs reveal a long arc of nebulosity curving up the east side Orion. This is called "Barnard's Loop" which is extremely difficult to discern visually almost regardless of aperture. Remarkably it has been seen with the naked eye (initially by accident!) from dark sites using O III or UHC filters. The "Loop" is a faint ring of hot gas some 14^o by 10^o with the western part of the ring being less distinct. The "ring" may be due to radiation pressure from the hot young stars in the region of Orion's belt/sword acting on interstellar material. A less favoured school of thought is that it may be a supernova remnant.

Taurus (Tau).

Lambda (λ) vs. Eclipsing binary, range 3.4 to 4.1, period 3.95d.

Phi (ψ) ds. 5.0/8.4 sep 52.1". Fine contrasting deep yellow primary with blue companion.

Σ 495 (Struve) ds. 6.0/8.8 sep 3.8". Fine pair of yellow stars.

Σ 742 ds. 7.1/7.5 sep 4". Fine pair in the same wide field as the Crab Nebula (M1) to the E. Often overlooked when locating M1.

47 Tauri ds. 4.9/7.4 sep 1.1". Very close pair of yellow stars.

T vs. Irregular variable, range 9.4 to 13.5, period erratic.

RV vs. Irregular variable, range 9.5 to 13, period 79d.

RR vs. Irregular variable, range 9.9 to 13, period "chaotic".

M45 - "The Pleiades" or "Seven Sisters" oc. Probably the most famous star cluster. Test your eyesight from a dark site by counting the number of naked eye stars that are visible. Seven should readily be seen. Keen vision will lead you into double figures. A test for moderate apertures is the nebulosity around some of the other brighter stars of the group, especially Merope. Nebula/UHC filters will help.

The Hyades oc. Another naked eye star cluster although more spread out than the Pleiades. The first magnitude star Aldebaran (0.9) is not a member of the cluster and appears brighter as it is closer to us, 21 parsecs - about halfway to the clusters centre. Aldebaran has a faint companion (13.4) separation 30.4". Spotting the companion is not easy even in moderate telescopes.

NGC1514 (10.0) pn. Not an easy object in small instruments. An OIII filter will help.

NGC1554/1555 is a faint variable reflection nebula illuminated by the very young star T Tauri (see above).

NGC1647 (6.4) oc. Rich cluster.

NGC1746 (6.1) oc. Poorer than 1647

NGC1808/1817 (7.0/7.7) oc. a pleasant "double cluster" when observed at low powers.

NGC1952 (**M1**)(8.5) snr. "The Crab Nebula". One of the most studied astronomical objects in recent decades. A "new star" appeared in 1054 and over a few months faded from view. Centuries later the faint oval patch was discovered by Dr John Bevis in 1731 and independently by Charles Messier on 12th September 1758 while observing the comet of that year. Its true nature was not recognized until much later. In 1942 astronomers had speculated that a peculiar star in the nebula might be a neutron star. In 1968 radio astronomers discovered that the central star was a pulsar spinning at 30 times a second i.e. the period of the pulses is 33 milliseconds. The "Crab" is also a strong emitter of X-rays.

Happy and Successful Observing!

P.V.H.